

Abdominal Pain

Abdominal (belly) pain can be caused by many things. Your caregiver decides the seriousness of your pain by an examination and possibly blood tests and x-rays. Many cases can be observed and treated at home. Most abdominal pain in children is functional. This means it is **not** caused by a disease and will probably improve without treatment. However, in many cases, more time must pass before a clear cause of the pain can be found. Before that point, it may not be known if you need more testing, or if hospitalization or surgery is needed.

HOME CARE INSTRUCTIONS:

- **Do not** take or give laxatives unless directed by your caregiver.
- Take pain medication only if ordered by your caregiver.
- Only take over-the-counter or prescription medicines for pain, discomfort or fever as directed by your caregiver.
- Try a clear liquid diet - broth, tea, or water as ordered by your caregiver. Slowly move to a bland diet as tolerated.

SEEK IMMEDIATE MEDICAL ATTENTION IF:

- The pain does not go away.
- An oral temperature above 102° F (38.9° C) develops or as directed by your caregiver.
- Repeated vomiting occurs.
- The pain is felt only in portions of the abdomen. The right side could possibly be appendicitis. In an adult, the left lower portion of the abdomen could be colitis or diverticulitis.
- Blood is being passed in stools (bright red or black tarry stools).

MAKE SURE YOU:

- Understand these instructions.
- Will watch your condition.
- Will get help right away if you are not doing well or get worse.

